



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

cooling of the mass." The original body of magma was leucocratic, so that the alumina and alkalis with silica enough for the formation of feldspar and feldspathoids, crystallized first, and hence in the periphery of the mass.

Densities of Liquid and Solid Rock Magmas.—In view of the fact that a recently proposed theory of volcanic action accounts for the eruption of lava as the consequence of the expansion of liquid magma in its passage to the solid state, a paper by Doelter,¹ in which this author discusses the densities of liquid and solid magmas, becomes of great interest. By means of a series of experiments, in the course of which fragments of known density were allowed to float or sink in molten magmas, Doelter obtained a series of results which are embodied in the following table, the figures indicating specific gravities:

	NATURAL SUBSTANCE.	MOLTEN.	GLASSY.	CRYSTALLINE.
Melanite. . . .	3.75	3.55-3.6	3.55-3.6	3.65-3.7
Augite	3.3	2.92	2.92-2.95	3.2-3.25
Limburgite . . .	2.83	2.55-2.57	2.55-2.57	2.75-2.78
Lava (Ætna) . .	2.83	2.58-2.74	2.71-2.75	2.81-2.83
Lava (Vesuvius) .	2.84	2.68-2.74	2.69-2.75	2.77-2.81
Nephelinite . . .	2.74	2.70-2.75	2.686	2.72-2.75
Leucitite	2.83	2.60-2.68	2.68-2.72	2.75-2.79

The Laccolite of Shefford Mountain.²—Shefford Mountain is the easternmost of the series of nine hills of igneous material that rise above the St. Lawrence valley in the neighborhood of Montreal. It is thought by Dresser to be an old laccolite in Lower Cambrian and Trenton sediments. Its material consists of essexite, nordmarkite, and pulaskite, the first two of which possess almost the typical character of these rocks. Associated with these are dark-colored dykes of a camptonitic type, and others of a theralitic type, and light-colored ones of trachyte and bostonite. The latter are the younger. All the rocks are thought to be differentiated products of a single magma. The primary magma, according to this view, had nearly the composition of pulaskite. Excluding the dykes the first differentiate was the basic essexite, the second was the acid nordmarkite, and the third the intermediate pulaskite. The analyses of the essexite (I),

¹ *Neues Jahrb. of Univ.*, etc., Bd. ii (1901), p. 141.

² *Amer. Geologist*, vol. xxvii (Oct., 1901), p. 205.

the nordmarkite (II), the pulaskite (III), and the mean between the first two (IV) follow:

SiO ₂	TiO ₂	Al ₂ O ₃	Fe ₂ O ₃	FeO	MnO	CaO	BaO	MgO	K ₂ O	Na ₂ O	P ₂ O ₅	CO ₂	SO ₃	Cl	H ₂ O	Total
53.15	1.52	17.64	3.10	4.65	.46	5.66	.13	2.94	3.10	5.00	.65	.39	.28	.07	1.10	= 99.86
65.43	.16	16.96	1.55	1.53	.40	1.36		.22	5.36	5.95	.02		.06	.04	.82	= 99.84
59.96	.66	19.12	1.85	1.73	.49	2.24	.12	.65	4.91	6.98	.14		.08	.14	1.10	= 100.17
59.29	.84	17.30	2.32	3.09	.43	3.51	.07	1.58	4.23	5.47	.34	.20	.17	.06	.96	= 99.85